HUMAN MOVEMENT 2017; 18 (1): 11-18



THE RELATIONSHIP BETWEEN STRENGTH MEASUREMENTS AND ANTHROPOMETRIC INDICATORS (BMI AND SKINFOLD THICKNESS) IN ELLISRAS RURAL ADOLESCENTS AGED 9-15 YEARS: **ELLISRAS LONGITUDINAL STUDY**

original paper doi: 10.1515/humo-2017-0003

MBELEGE R. NKWANA, KOTSEDI D. MONYEKI, MOLOKO MATSHIPI, MACHOENE D. SEKGALA, NTHAI E. RAMOSHABA, TSHEPHANG M.J. MASHIANE

Department of Physiology and Environmental Health, University of Limpopo, Sovenga, South Africa

ABSTRACT

Purpose. Handgrip strength and arm hang have been recognized as predictors of muscle strength and presented as biomarkers for important health outcomes and overall fitness of an individual. The aim of the study was to investigate the relationship between arm hang and handgrip strength with BMI and skinfold thickness.

Method. The total of 769 children (391 boys and 378 girls) aged 9-15 years underwent a series of anthropometric and strength measurements with the use of standard procedures. Linear regression was applied to assess the relationship of arm hang and handgrip strengths with body mass index and skinfold thickness.

Results. The prevalence of undernutrition and low strength measurements was high (1.7–85%) while the prevalence of overweight and obesity was low (1.5-4.2%) in the Ellisras children aged 9–15 years. Boys showed a significantly higher (p < 0.05) mean arm hang (9.6-13.1 kg) than girls (4.0-5.1 kg) aged 11–15 years. There was a significant (p < 0.05) association between low strength and underweight among Ellisras rural children when unadjusted (OR = 0.650; 95% CI = 0.487-0.867) and adjusted (OR = 0.650; 95% CI = 0.489–0.868) for age and gender.

Conclusions. The prevalence of undernutrition and poor strength was high. An association was observed between arm hang and anthropometric indicators while handgrip showed no significant association with anthropometric indicators. Further studies are needed to investigate the association between strength and undernutrition over time.

Key words: undernutrition, upper body strength, rural South African children

Introduction

Malnutrition is a risk factor for ill health and contributes greatly to the burden of disease in low- to middleincome countries [1]. It increases the risk of cardiovascular diseases among children and adults in the developed and developing countries [2-5]. Body mass index (BMI) has been used widely as an indicator for malnutrition, although it has been recently argued that BMI reflects also the relative leg length, body frame size, and fat free mass in addition to fatness [6, 7].

Koley and Kaur [8] suggest that skinfold thickness, arm hang, and handgrip strength can be used in addition to BMI to screen malnutrition. Skinfold thickness has been reported as a good indicator for malnutrition as it can measure the distribution of subcutaneous adipose fat [9]. Poor strength was proved to be associated with underweight, while good strength was bound with overweight in an individual [10].

Koley and Singh [11] reported a significant association between handgrip strength and BMI among Amritsar youth aged 6-25 years, while Freedman et al. [12] observed a significant association between handgrip strength and skinfold thickness among children and adolescents aged 5-18 years from the USA. Monyeki et al. [13, 14] described an inverse relationship of bent arm hang with weight for age z-score, sum of four skinfolds, and fat free mass while the prevalence of undernutrition was high amongst Ellisras rural children. The association of BMI with skinfold thickness, handgrip, and arm hang strength has received little attention in rural South Africans. Therefore, the aim of this study was to investigate 1) the prevalence of malnutrition and muscle strength, 2) the relationship of arm hang and handgrip strength with BMI and skinfold thickness, 3) the risk of developing undernutrition among Ellisras rural children aged 9-15 years.

Received: July 30, 2016 Acepted for publication: February 14, 2017

Correspondence address: Kotsedi D. Monyeki, Department of Physiology and Environmental Health, University of Limpopo, Private Bag X 1106, Sovenga 0727, South Africa, e-mail: dmonyeki@yahoo.com

Citation: Nkwana MR, Monyeki KD, Matshipi M, Sekgala MD, Ramoshaba NE, Mashiane TMJ. The relationship between strength measurements and anthropometric indicators (BMI and skinfold thickness) in ellisras rural adolescents aged 9-15 years: Ellisras longitudinal study. Hum Mov. 2017;18(1):11-18; doi: 10.1515/humo-2017-0003.

Material and methods

Geographical area

Ellisras is a deep rural area situated within the north western area of the Limpopo province, South Africa. The population is about 50,000 people, residing in 42 settlements [15]. These villages are approximately 70 km away from the Ellisras town (231 40S 271 44W), now known as Lephalale, adjacent to the Botswana border. The Iscor coal mine and Matimba electricity power station are the major sources of employment for many of the Ellisras residents, whereas the remaining workforce is involved in subsistence farming and cattle rearing, and a minority – in education and the civil service. Unemployment, poverty, and low life expectancy seem to play a significant role in the rural South African population, to which the Ellisras rural area people are not an exception [16].

Sample and research design

Details of the Ellisras Longitudinal Study design and sampling have been reported elsewhere [17, 18]. The total of 769 subjects (391 boys and 378 girls) aged 9–15 years who completed all the anthropometric and strength measurements were included in the analysis. The Ethics Committee of the University of Limpopo granted ethical approval prior to the survey, and the participants' parents or guardians provided their written informed consents.

Anthropometric measurement

All the children underwent a series of anthropometric measurements according to the standard procedures recommended by the International Society of the Advancement of Kinanthropometry (ISAK) [19]. Weight was measured on an electronic scales and rounded to the nearest 0.1 kg, and a Martin anthropometric was used to define height rounded to the nearest 0.1 cm. Skinfolds (suprailiac, subscapular, triceps, biceps) were measured three times with the use of a Slim Guide skinfold calliper, and the values were rounded to the nearest 0.1 mm. BMI was defined as weight (kg)/height (m)².

Arm hang and handgrip strength measurements

Arm hang and handgrip strength were measured with standard procedures described by the European tests of physical fitness [20]. Arm hang was determined with the use of a horizontal bar, a chair, and a stopwatch. The participant grabbed the bar with an overhand and gripped so that the palms were facing away, then the participant raised their body off the floor/chair so that the chin went above the bar, the elbows were flexed, and the chest was near the bar. The stopwatch was started immediately when the participant was at the hanging position. The position was maintained for as long as possible. Handgrip was measured in both hands with a dynamometer. The participant's elbow was flexed at the 90° angle, with the forearm parallel to the floor. The dynamometer was maximally squeezed for a 3-second count while, simultaneously lowering the arm to full extension. The participants practiced this procedure once per hand, after which measurements were recorded and rounded to the nearest 0.5 kg. The instrumental precision error was 1.0% [21].

Statistical analysis

Descriptive statistics were expressed for handgrip, arm hang strengths, BMI, and skinfold thickness. The t-test was used to compare the significant differences between genders by age group. Subscapular and triceps skinfolds of the Ellisras rural children were compared with the National Health and Nutrition Examination Survey III (NHANES) reference population. The international cut-off points for underweight (grade one, two, and three) by sex for exact ages defined to pass through the BMI of 16, 17, and 18 kg/m² were used [22]. The children were categorized as normal and over fatness with the use of the sum of four skinfolds (subscapular, triceps, biceps, and suprailiac) above 90th percentiles [23]. The following cut off points for strength measurements were used: scores below 25th percentile were poor, above the 25th percentile but below the 50th percentile – minimal fitness, above the 50th percentile but below the 75th percentile – good strength, and scores at or above the 75th percentile represented excellent strength. Chi-squared tests were applied to compare sets of nominal data that had larger frequency counts, whereas Fisher's exact test was used when frequency cells were small (less than five or ten) between genders.

Pearson correlation coefficient was performed to determine the relationship between handgrip, arm hang strength, BMI, and skinfold thickness by gender. Linear regression coefficient analysis was used to assess the relationship between handgrip, arm hang strengths, BMI, and skinfolds thickness, both unadjusted and adjusted for age and gender. Logistic regression allowed to estimate the association between strength measurements and the odds of incident malnutrition (underweight) while adjusting for covariate known to be associated with malnutrition (age, gender). All the data were analysed with a statistical package for social science (SPSS), version 23. The statistically significant difference was assumed at p < 0.05.

Results

Figures 1 and 2 show a comparison between triceps and subscapular skinfolds of the NHANES III reference population and Ellisras children aged 9–15 years. The NHANES III reference population showed higher M.R. Nkwana et al., Strength measurements and anthropometric indicators



Figure 1. Comparison of the National Health and Nutrition Examination Survey III (NHANES) reference population and Ellisras (ELS) rural children aged 9–15 years with reference to triceps skinfold thickness

triceps and subscapular skinfolds than the Ellisras rural children throughout the age range (9–15 years).

Table 1 shows the descriptive statistics of absolute body size, handgrips (left and right), and arm hang strength of Ellisras rural children aged 9–15 years. Girls aged 11–13 years represented significantly higher (p < 0.05) mean BMI values (15.5–16.5 kg/m²) compared with boys (14.7–15.4 kg/m²) of the same age. In turn, girls aged 11–12 years had significantly higher (p < 0.05) mean



Figure 2. Comparison of the National Health and Nutrition Examination Survey III (NHANES) reference population and Ellisras (ELS) rural children aged 9–15 years with reference to subscapular skinfold thickness

sum of four skinfolds (26.3–29.6 cm) than boys (21.5–22.1 cm) of the same age. In boys, a significantly higher (p < 0.05) mean arm hang was observed (9.6–13 s) than in girls (4.0–5.1 s) at the age of 11–15 years.

In Table 2, the prevalence of strength measurements among Ellisras children aged 9–15 years is presented. The prevalence of poor strength (arm hang and hand grip) was higher among girls (41–73%) than in boys (20–55.3%) at the age of 12–15 years.

Table 1. The descriptive statistics of absolute body size, hand grips (left and right), and arm hang strengths in Ellisras rural children aged 9–15 years (N = 769)

			Right h (k	andgrip g)) Left ha	andgrip (g)	Arm (hang s)	Bl (kg/	MI (m2)	Bic (m	eps m)	Supr (m	ailiac 1m)	Sur skinfole (biceps, supra subsca	n of ds (mm) triceps, uiliac, upular)
Age (years)	Boys (<i>n</i> = 391)	Girls (<i>n</i> = 378)	boys m (SD)	girls m (SD)	boys m (SD)	girls m (SD)	boys m (SD)	girls m (SD)	boys m (SD)	girls m (SD)	boys m (SD)	girls m (SD)	boys m (SD)	girls m (SD)	boys m (SD)	girls m (SD)
9	35	24	15.3 (6.1)	13.6 (5.9)	13.9 (5.8)	12.1 (5.2)	14.5** (11.7)	5.2** (4.3)	14.4 (1.3)	14.4 (2.3)	3.9* (0.9)	4.5* (2.4)	4.5 (1.2)	4.4 (2.3)	20.3* (4.4)	20.7* (9.2)
10	47	52	15.3 (6.2)	12.9 (5.7)	14.1 (6.2)	12.0 (5.4)	7.2 (5.9)	4.6 (4.4)	14.6 (1.3)	14.6 (1.4)	4.1* (1.1)	4.7* (1.4)	4.6 (1.4)	4.9 (1.8)	21.0 (4.9)	23.2 (5.8)
11	68	55	13.1 (5.2)	13.7 (6.0)	12.0 (4.9)	12.4 (5.9)	10.0* (8.1)	5.1* (4.3)	14.7* (1.4)	15.5* (2.1)	4.2* (1.5)	5.4* (2.0)	5.1 (1.9)	5.7 (1.8)	22.1* (6.7)	26.3* (8.5)
12	69	76	15.1 (6.2)	13.5 (5.8)	14.3 (5.7)	12.4 (6.9)	9.6* (8.3)	4.0** (3.5)	15.1* (1.4)	16.4* (2.5)	4.0* (1.0)	5.6* (2.5)	4.8* (1.5)	7.1* (3.9)	21.5* (4.9)	29.6* (12.4)
13	68	65	14.0 (5.0)	14.4 (6.9)	13.0 (5.0)	13.2 (6.7)	13.1** (10.6)	4.9** (7.1)	15.4* (1.5)	16.5* (2.6)	4.4 (1.6)	5.5 (2.0)	5.6* (2.7)	7.0* (3.2)	23.8 (8.1)	29.6 (10.3)
14	61	60	15.0 (6.6)	14.3 (5.3)	13.9 (6.6)	13.0 (5.3)	11.6** (9.6)	4.6** (3.9)	16.2 (1.7)	17.2 (1.7)	4.6 (1.7)	5.8 (1.9)	6.1 (3.0)	9.2 (4.0)	26.0 (9.3)	34.5 (11.3)
15	43	46	13.7 (6.1)	13.7 (5.1)	12.8 (5.6)	12.9 (5.3)	12.4* (7.6)	5.1* (5.0)	16.9 (1.6)	17.8 (1.2)	4.3 (1.7)	5.6 (1.3)	6.0 (2.0)	7.7 (2.5)	24.4 (7.3)	32.1 (8.7)

N – number of population, m – mean, SD – standard deviation, * p < 0.05, ** p < 0.001

M.R. Nkwana et al., Strength measurements and anthropometric indicators

					°	•					
	Ι	Left handgrij	0	R	ight handgri	ip	Arm hang				
Age (years)	minimal strength % (<i>n</i>)	good % (<i>n</i>)	excellent % (<i>n</i>)	minimal strength % (n)	good % (<i>n</i>)	excellent % (<i>n</i>)	minimal strength % (n)	good % (<i>n</i>)	excellent % (<i>n</i>)		
Boys											
9	40.0 (14)	28.6 (10)	31.4 (11)	37.1 (13)	20.0 (7)	42.9 (15)	20 (7)	34.3 (12)	45.7 (16)		
10	40.5 (19)	29.8 (14)	29.8 (14)	31.9 (15)	36.2 (17)	31.9 (15)	55.3 (26)	19.1 (9)	25.5 (12)		
11	50 (34)	30.9 (21)	19.1 (13)	48.5 (33)	32.4 (22)	19.1 (13)	25 (17)	32 (26)	36.8 (25)		
12	40.6 (28)*	27.5 (19)	31.9 (22)	39.1 (27)*	30.4 (21)	30.4 (21)	43.5 (30)	21.7 (15)	34.8 (24)		
13	45.6 (31)	30.9 (21)	23.5 (16)	41.2 (28)*	35.3 (24)	23.5 (16)	26.5 (18)*	23.5 (16)	50 (34)		
14	45.9 (28)*	21.3 (13)	32.8 (20)	41.1 (25)	26.2 (16)	62.8 (20)	29.5 (18)*	29.5 (18)	41 (25)		
15	46.5 (20)	25.6 (11)	22.9 (12)	51.2 (22)	20.9 (9)	27.9 (12)	25.6 (11)*	16.3 (7)	58.1 (24)		
				Gi	rls						
9	45.8 (11)	12.5 (3)	41.7 (10)	50 (12)	33.3 (8)	16.7 (4)	58.3 (14)	29.2 (7)	12.5 (3)		
10	55.8 (29)	23.1 (12)	21.2 (11)	53.8 (28)	28.8 (15)	17.3 (9)	73.1 (38)	21.2 (11)	5.8 (3)		
11	49.1 (27)	30.9 (17)	20.0 (11)	47.3 (26)	29.1 (16)	23.6 (13)	67.3 (37)	18.2 (10)	14.5 (8)		
12	55.3 (42)	22.4 (17)	22.4 (17)	51.3 (39)*	27.6 (21)	21.1 (16)	72.4 (55)	23.7 (18)	3.9 (3)		
13	62.3 (34)	20.0 (13)	22.4 (18)	50.8 (33)*	21.5 (14)	27.7 (18)	69.2 (45)*	24.6 (16)	6.2 (4)		
14	50 (30)*	23.3 (14)	26.7 (16)	41.0 (25)	36.7 (22)	21.7 (13)	65.0 (39)*	28.3 (17)	6.7 (4)		
15	50 (23)	26.1 (12)	23.9 (11)	50.0 (23)	28.3 (13)	21.7 (10)	63.0 (29)*	21.7 (10)	15.2 (7)		

Table 2. The prevalence of strength measurements (left and right handgrip and arm hang)	
in Ellisras children aged 9–15 years	

* *p* < 0.05

Table 3. The prevalence of malnutrition (severe, moderate, and mild underweight, overweight, obese, and over fatness) as established by the body mass index and sum of four skinfolds (triceps, biceps, suprailiac, and subscapular) among Ellisras rural children aged 9–15 years

Age group	(N = 769)		Severe underweight		Moderate underweight		Mild underweight		Overweight		Obese		Over fatness	
group (years)	boys (<i>n</i> = 391)	girls (<i>n</i> = 378)	boys % (<i>n</i>)	girls % (<i>n</i>)	boys % (<i>n</i>)	girls % (<i>n</i>)	boys % (<i>n</i>)	girls % (<i>n</i>)	boys % (<i>n</i>)	girls % (<i>n</i>)	boys % (<i>n</i>)	girls % (n)	Over fat	girls % (<i>n</i>)
9	35	24	2.9 (1)	12.5 (3)	22.9 (8)	16.7 (4)	28.6 (10)	33.3 (8)	-	4.2 (1)	_	_	8.6 (3)	8.3 (2)
10	47	52	12.8 (6)	3.8 (2)	-	11.5 (6)	44.7 (21)	36.5 (19)	-	-	-	-	8.5 (4)	9.6 (5)
11	68	55	8.8 (6)	5.5 (3)	14.7 (10)	9.1 (5)	36.8 (25)	32.7 (18)	1.5 (1)	3.6 (2)	-	-	8.8 (6)	9.1 (5)
12	69	76	8.7 (6)	7.9 (6)	10.1 (7)	7.9 (6)	42.0 (29)	23.7 (18)	-	2.6 (2)	-	-	8.7 (6)	9.2 (7)
13	68	65	10.3 (7)	12.3 (8)	10.3 (7)	16.9 (11)	50.0 (34)	29.2 (19)	-	3.1 (2)	-	-	8.8 (6)	9.2 (6)
14	61	60	13.1 (8)	1.7 (1)	14.8 (9)	13.3 (8)	23.0 (14)	85.0 (51)	-	-	-	-	9.8 (6)	10.0 (6)
15	43	46	9.3 (4)	4.3 (2)	16.3 (7)	6.5 (3)	25.6 (11)	30.4 (14)	-	-	-	-	9.3 (4)	8.7 (4)

Table 3 illustrates the prevalence of malnutrition in Ellisras children aged 9–15 years. The majority of Ellisras rural children were underweight (1.7-85%), and few were overweight (1.5-4.2%). The prevalence of over fatness ranged from 8.3 to 10.0%, and the difference between genders was insignificant.

The Pearson correlation between handgrip strength, arm hang strength, and other anthropometric indicators can be seen in Table 4. There was a significant (p < 0.001) negative association (r^2 ranged from -0.24 to -0.13) between arm hang strength and all other anthropometric indicators.

Table 5 presents the linear regression analysis. A significant (p < 0.05) positive association was observed

between right handgrip and triceps ($\beta = 0.191$; 95% CI = 0.026–0.356), unadjusted. Moreover, right handgrip showed a significant positive association with biceps ($\beta = 0.168$; 95% CI = 0.00–0.337), unadjusted. When adjusted for age and gender, left handgrip was negatively associated with biceps ($\beta = -0.186$; 95% CI = from -0.353 to -0.018) among Ellisras rural children.

The logistics regression of the association between poor strengths and nutrition status among Ellisras rural children aged 9–15 years is illustrated in Table 6. There was a significant (p < 0.05) association between poor strength and underweight, both unadjusted (OR = 0.60; 95% CI = 0.43–0.85) and adjusted (OR = 0.60; 95% CI = 0.43–0.84) for age and gender among Ellisras rural children.

Variable	Right h	andgrip	Left ha	ndgrip	Arm hang		
variable	boys	girls	boys	girls	boys	girls	
BMI	-0.02	-0.03	-0.01	-0.06	-0.13*	-0.21**	
Triceps	0.02	-0.03	0.04	-0.06	-0.23**	-0.21**	
Biceps	-0.03	-0.02	-0.04	-0.07	-0.24**	-0.20**	
Subscapular	-0.03	-0.04	-0.00	-0.07	-0.14**	-0.19**	
Suprailiac	-0.05	-0.05	-0.04	-0.08	-0.19**	-0.18**	
Sum of skinfold	-0.01	-0.02	0.01	-0.06	-0.20**	-0.18**	

Table 4. The Pearson moment correlation coefficient of absolute body size and strength measurements of Ellisras children aged 9–15 years

BMI – body mass index, * *p* < 0.05, ** *p* < 0.001

Table 5. Linear regression coefficients, *p* value, and 95% confidence intervals for the association of skinfolds and body mass index with handgrips and bent arm hang among Elliras children aged 9–15 years

		Unadju	sted	Adjusted (age and gender)					
variable	β	<i>p</i> value	95% CI		β	<i>p</i> value	95% CI		
Right handgrip									
BMI	-0.10	0.344	-0.30	0.10	-0.07	0.537	-0.30	0.16	
Triceps	0.19	0.023*	0.03	0.36	-0.04	0.425	-0.15	0.06	
Biceps	-0.01	0.943	-0.16	0.15	0.17	0.050*	0.00	0.34	
Subscapular	-0.05	0.416	-0.16	0.07	-0.00	0.964	-0.15	0.15	
Suprailiac	0.01	0.720	-0.02	0.03	-0.05	0.399	-0.16	0.07	
Sum of skinfolds	-0.03	0.181	-0.07	0.01	-0.02	0.371	-0.07	0.03	
Left handgrip									
BMI	-0.03	0.644	-0.15	0.09	-0.06	0.345	-0.19	0.07	
Triceps	0.08	0.157	-0.03	0.18	0.08	0.150	-0.03	0.19	
Biceps	-0.22	0.009*	-0.38	-0.05	-0.19	0.030*	-0.35	-0.02	
Subscapular	-0.00	0.988	-0.15	0.15	-0.01	0.891	-0.16	0.14	
Suprailiac	0.01	0.801	-0.10	0.13	0.01	0.895	-0.11	0.12	
Sum of skinfolds	-0.039	0.075	-0.083	0.004	-0.036	0.144	-0.083	0.012	
Arm hang									
BMI	-0.81	0.000	-0.08	-0.55	-0.70	0.000	-0.98	-0.35	
Triceps	-0.69	0.000	-0.86	-0.53	-0.52	0.000	-0.69	-0.35	
Biceps	-1.29	0.000	-1.59	-0.99	-0.86	0.000	-1.16	-0.57	
Subscapular	-0.81	0.000	-1.03	-0.57	-0.55	0.000	-0.78	-0.31	
Suprailiac	-0.68	0.000	-0.87	-0.49	-0.51	0.000	-0.71	-0.32	
Sum of skinfolds	-0.24	0.000	-0.31	-0.19	-0.18	0.000	-0.24	-0.12	

CI – confidence interval, BMI – body mass index, * p < 0.05

Table 6. Odds ratio and 95% confidence interval for the association of poor strengths and undernutrition
among Ellisras rural children aged 7–11 years

W		Unadjusted		Adjusted for age and gender					
variable	OR	<i>p</i> value	95%	6 CI	OR	<i>p</i> value	nd gender 95% * 0.43 0.81	6 CI	
Poor strength									
Underweight	0.60	0.003**	0.43	0.84	0.60	0.004**	0.43	0.85	
Over fatness	1.50	0.195	0.81	2.75	1.52	0.190	0.81	02.84	

CI – confidence interval, ** p < 0.001

Discussion

The aim of the study was to investigate the association between handgrip, arm hang strength, BMI, and skinfold thickness of rural Ellisras children aged 9–15 years. The strength test was used to evaluate the performance of hand muscle by measuring the maximum grip force that could be executed by one muscular contraction [24]. There were significant associations between skinfold, BMI, and arm hang strength in the population. No significant associations were recorded between anthropometric indicators and grip strength measurements.

Generally, boys showed higher muscular strength than girls (left and right handgrip and arm hang) [2], though in another study [25] no significant difference between genders was observed. The grip strength was reported to be higher in the dominant hand but no such significant differences between genders could have been documented [26] as in the current study, in which boys had significantly higher handgrip and arm hang strength than girls. Similar results were reported among European and Indian children aged 6–17 years [7, 27, 28].

The prevalence of underweight and low fatness was high among Ellisras rural children aged 9–15 years. Furthermore, NHANES III reference children presented higher subscapular and triceps measurements than Ellisras rural children. Similar results were earlier reported with regard to poor nutritional status among rural South African children [29–32]. The prevalence of poor muscle strength was high in the current study. Furthermore, Monyeki et al. [13] observed low physical activity in the sample as compared with children studied in other parts of the world. Poor muscle strength was reported to be significantly associated with lower body weight or undernutrition, presence of chronic diseases, and physical inactivity [33, 34], while good strength could be an indicator of better childhood and early life nutrition [35].

In the present study, arm hang strength was negatively associated with BMI and skinfold thickness. Handgrip strength did not show any significant association with other anthropometric indicators. The study results remain in contrast with previous studies which reported a significant positive association between hand grip strength and skinfold thickness among children and adolescents [12, 35]. A decrease in muscle mass reduces physical strength, with low energy available to be used owing to low fatness among Ellisras rural children.

Our findings suggest that underweight children with low body fatness were more likely to develop low strength than children with normal body fatness. The results were similar to a previous study by Must et al. [36], who reported that underweight was significantly associated with poor strength among Indian youth. Physiologically, this could be due to poor dietary or energy intake that leads to underweight and poor strength [5, 37]. Furthermore, thinner people often have poor strength, higher illness prevalence, and greater mortality than those with normal body weight [35]. Poor muscle strength could be a risk for diseases later in life.

The major limitation of the study is that our sample only included children from rural areas of South Africa, dominated by an indigenous knowledge system [38]; as a result; extrapolation of the outcomes to urban areas should be made with caution. In our study, we did not consider the education level of the subjects, which has been reported to be associated with poor muscular strength [39]. Maturation stage, physical fitness, physical activity, and the family socioeconomic level of the participants could not be included in the analysis.

Conclusions

The prevalence of undernutrition and low strength measurements were high in the Ellisras rural children. There was a significant association between arm hang and other anthropometric indicators, while handgrip showed no significant association with other anthropometric indicators. Further studies are needed to investigate the association between strength and undernutrition overtime.

Acknowledgements

The authors would like to thank R. Frankhuisen and W. Kars (4th year Vrije Medical University students) for the data collection in the Ellisras Longitudinal Study. The financial support received from Vrije University, Amsterdam, the Netherlands; University of Limpopo, South Africa; and the National Research Foundation of South Africa for the Ellisras Longitudinal Study is also acknowledged with gratitude. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the authors and therefore the above mentioned funding sources do not accept any liability in regard thereto. The authors are indebted to ELS administrators (P.S. Seleka, T.T. Makata, W. Makata, S. Seleka) for providing technical support in the preparation of the manuscript. M.S. Monyeki and M.J. Malatji (Makgoka Secondary School, English educators, Limpopo province) are thankfully acknowledged for editing the manuscript.

Funding

The work was supported by South African National Research Foundation (grant No. URD2002050400168), University of Limpopo (grant No. 1404), Vu Medical University, Amsterdam (grant: UNIN Health Project under VUA Foundation funds).

References

- 1. World Health Organization, The world health report 2002 reducing risks, promoting healthy life. Geneva: WHO; 2002.
- 2. Bragagnolo R, Caporossi FS, Dock-Nascimento DB, de Aguilar-Nascimento JE. Handgrip strength and adductor pollicis muscle thickness as predictors of postoperative complications after major operations of the gastrointestinal tract. Eur E J Clin Nutr Metab. 2011;6(1):21–26; doi: 10.1016/j.eclnm.2010.11.001.
- 3. Garcia MF, Wazlawik E, Moreno YMF, Führ LM, González-Chica DA. Diagnostic accuracy of handgrip strength in

the assessment of malnutrition in hemodialyzed patients. Eur E J Clin Nutr Metab. 2013;8(4):181–186, doi: 10.1016/ j.clnme.2013.06.003.

- 4. Kemper HCG (ed.). Amsterdam growth and health longitudinal study. A 23-year follow-up from teenager to adult about lifestyle and health. Basil: Karger; 2004.
- 5. Monyeki KD, Kemper HCG. The risk factors for elevated blood pressure and how to address cardiovascular risk factors: a review in paediatric populations. J Hum Hypertens. 2008;22(7):450–459; doi: 10.1038/jhh.2008.21.
- 6. Gale CR, Martyn CN, Cooper C, Sayer AA. Grip strength, body composition, and mortality. Int J Epidemiol. 2007; 36(1):228–235; doi: 10.1093/ije/dyl224.
- 7. Gandhi M, Koley S, Sandhu JS. Association between anthropometry characteristics and physical strength in school going children of Amritsar. Anthropologist. 2010; 12(1):35–39.
- 8. Koley S, Kaur N. A study on handgrip strength and some anthropometric variables in younger and older female laborers of Jalandhar, Punjab, India. Int J Biol Anthrop. 2010;3(2).
- 9. Sengupta P, Chaudhuri P, Bhattacharya K. Screening obesity by direct and derived anthropometric indices with evaluation of physical efficiency among female college students of Kolkata. Ann Medical Health Sci Res. 2013; 3(4):517–522; doi: 10.4103/2141-9248.122066.
- McRae MP. Male and female differences invariability with estimating body fat composition using skinfold calipers. J Chiropr Med. 2010;9(4);157–161; doi: 10.1016/j.jcm. 2010.07.002.
- 11. Koley S, Singh AP. An association of dominant hand grip strength with some anthropometric variables in Indian collegiate population. Anthropol Anz. 2009;67(1):21–28, doi: 10.1127/0003-5548/2009/0003.
- 12. Freedman DS, Wang J, Ogden CL, Thornton JC, Mei Z, Pierson RN, et al. The prediction of body fatness by BMI and skinfold thicknesses among children and adolescents. Ann Hum Biol. 2007;34(2):183–194; doi: 10.1080/030 14460601116860.
- 13. Monyeki MA, Koppes LLJ, Monyeki KD, Kemper HCG, Twisk JWR. Longitudinal relationships between nutritional status, body composition, and physical fitness in rural children of South Africa: the Ellisras longitudinal study. Am J Hum Biol. 2007;19(4):551–558; doi: 10.1002/ ajhb.20616.
- 14. Monyeki KD, Monyeki MA, Brits SJ, Kemper HCG, Makgae PJ. Development and tracking of body mass index from preschool age into adolescence in rural South African children: Ellisras longitudinal growth and health study. J Health Popul Nutr. 2008;26(4):405–417.
- 15. Sidiropoulos E, Jeffery A, Mackay S, Gallocher R, Forgey H, Chips C. South Africa survey 1995/1996. Johannesburg: South African Institute of Race and Relations; 1996.
- Statistics South Africa, Cause of death in South Africa 1997–2001: Advance release of recorded causes of death. Pretoria: Statistics South Africa; 2002.
- 17. Monyeki KD, Cameron N, Getz B. Growth and nutritional status of rural South African children 3–10 years old: The Ellisras growth study. Am J Hum Biol. 2000;12(1):42–49; doi: 10.1002/(SICI)1520-6300(200001/02)12:1<42::AID-AJHB6>3.0.CO;2-0.
- Monyeki KD, van Lenthe FJ, Steyn NP. Obesity: does it occur in African children in a rural community in South Africa? Int J Epidemiol. 1999;28(2):287–292; doi: 10.1093/ ije/28.2.287.

- Norton K, Olds T (eds.). Anthropometrica. A textbook of body measurement for sports and health courses. Sydney: University of New South Wales Press; 1996.
- 20. European Communities, Eurofit: European tests of physical fitness. Rome: Rome Council of Europe Committee for the development of sports; 1988.
- 21. Siengthai B, Kritz-Silverstein D, Barrett-Connor E. Handedness and cognitive function in older men and women: a comparison of methods. J Nutr Health Aging. 2008;12(9): 641–647, doi: 10.1007/BF03008275.
- 22. Cole TJ, Flegal KM, Nicholls D, Jackson AA. Body mass index cut offs to define thinness in children and adolescents: international survey. BMJ. 2007;335(7612):194; doi: 10.1136/bmj.39238.399444.55.
- 23. Twisk JW, Van Mechelen W, Kemper HCG, Post GB. The relation between "long-term exposure" to lifestyle during youth and young adulthood and risk factors for cardio-vascular disease at adult age. J Adolesc Health. 1997;20(4): 309–319; doi: 10.1016/S1054-139X(96)00183-8.
- 24. Massy-Westropp NM, Gill TK, Taylor AW, Bohannon RW, Hill CL. Hand grip strength: age and gender stratified normative data in a population-based study. BMC Res Notes. 2011;4:127; doi: 10.1186/1756-0500-4-127.
- 25. Link L, Lukens S, Bush MA. Spherical grip strength in children 3 to 6 years of age. Am J Occup Ther. 1995;49(4): 318–326; doi: 10.5014/ajot.49.4.318.
- 26. Incel NA, Ceceli E, Durukan PB, Erdem HR, Yorgancioglu ZR. Grip strength: effect of hand dominance. Singapore Med J. 2002;43(5):234–237.
- 27. Gulías-González R, Martínez-Vizcaíno V, García-Prieto JC, Díez-Fernández A, Olivas-Bravo A, Sánchez-López M. Excess of weight, but not underweight, is associated with poor physical fitness in children and adolescents from Castilla-La Mancha, Spain. Eur J Pediatr. 2014;173(6):727–735; doi: 10.1007/s00431-013-2233-y.
- Jürimäe J, Jürimäe T, Leppik A, Kums T. The influence of ghrelin, adiponectin, and leptin on bone mineral density in healthy postmenopausal women. J Bone Miner Metab. 2008;26(6):618–623; doi: 10.1007/s00774-008-0861-5.
- 29. Jinabhai CC, Reddy P, Taylor M, Monyeki D, Kamabaran N, Omardien R. Sex differences in under and over nutrition among school-going Black teenagers in South Africa: An uneven nutrition trajectory. Trop Med Int Health. 2007; 12(8):944–952; doi: 10.1111/j.1365-3156.2007.01861.x.
- 30. Kruger HS, Kruger A, Vorster HH, Jooste PL, Wolmarans P. Urbanization of Africans in the North West Province is associated with better micronutrient status: the transition and health during urbanization study in South Africa. Nutr Res. 2005;25(4):365–375; doi: 10.1016/j. nutres.2005.01.004.
- Labadarios D, Steyn N, Maunder E, MacIntyre U, Swart R. The National Food Consumption Survey (NFCS) – children aged 1–9 years, South Africa, 1999. South Afr J Clin Nutr. 2001;14(2):16.
- 32. Steyn K, Fourie J, Bradshaw D. The impact of chronic diseases of lifestyle and their major risk factors on mortality in South Africa. S Afr Med J. 1992;82(4):227–231.
- Artero EG, Lee DC, Ruiz JR, Sui X, Ortega FB, Church TS, et al. A prospective study of muscular strength and all-cause mortality in men with hypertension. J Am Coll Cardiol. 2011;57(18):1831–1837; doi: 10.1016/j.jacc.2010.12.025.
- Bartrina F. Theory and translator training. In: Tennent M. (ed.), Training for the new millennium: Pedagogies for translation and interpreting. Amsterdam, Philadelphia: John Benjamins Publishing Company, 2005; 177–189.

M.R. Nkwana et al., Strength measurements and anthropometric indicators

- 35. Rantanen T, Harris T, Leveille SG, Visser M, Foley D, Masaki K., et al. Muscle strength and body mass index as long-term predictors of mortality in initially healthy men. J Gerontol A Biol Sci Med Sc. 2000;55(3):168–173; doi: 10.1093/gerona/55.3.M168.
- 36. Must A, Dallal GE, Dietz WH. Reference data for obesity: 85th and 95th percentiles of body mass index (wt/ht2) and triceps skinfold thickness. Am J Clin Nutr. 1991;53(4): 839–846.
- 37. Van Den Ende C, Twisk JWR., Monyeki KD. The relationship between BMI and dietary intake of primary school children from a rural area of South Africa: The Ellisras longitudinal study. Am J Hum Biol. 2014;26(5):701–706; doi: 10.1002/ajhb.22585.
- Monyeki KD, Kemper HC, Makgae PJ. Development and tracking of central patterns of subcutaneous fat of rural South African youth: Ellisras longitudinal study. BMC Pediatr. 2009;9(1):74; doi: 10.1186/1471-2431-9-74.
- Rantanen T, Parkatti T, Heikkinen E. Muscle strength according to level of physical exercise and educational background in middle-aged women in Finland. Eur J Appl Physiol Occup Physiol. 1992;65(6):507–512; doi: 10.1007/ BF00602356.