



Effects of forward head posture on the centre of pressure sway during dual-task balancing with smartphone engagement

original paper

DOI: <https://doi.org/10.5114/hm/192231>

© Wrocław University of Health and Sport Sciences

WEERASAK TAPANYA^{ID}, NOPPHARATH SANGKARIT^{ID}

Department of Physical Therapy, School of Allied Health Sciences, University of Phayao, Phayao, Thailand

ABSTRACT

Purpose. Forward head posture (FHP) may affect postural stability and balance. This study examines how smartphone texting while balancing, as a dual task, influences the body's centre of pressure (CoP) during single-leg standing in individuals with FHP compared to those with normal posture.

Methods. The study utilized a randomized repeated measures design to assess single-leg standing balance and CoP across six conditions. Fifty participants, divided into two groups (25 with FHP and 25 with normal posture), underwent testing sequences involving non-smartphone usage (NU; single task), one-handed smartphone usage (1H; dual task), and two-handed smartphone usage (2H; dual task) on both soft and hard surfaces.

Results. The study found that individuals with FHP showed increased CoP sway compared to those with normal posture across all conditions. Both 1H and 2H increased sway compared to non-usage for both groups, with a notable difference in FHP individuals on soft surfaces, where 2H led to greater sway.

Conclusions. Individuals with FHP are at a higher risk of balance issues during both NU and smartphone use, particularly during 2H. These results highlight the importance of addressing the impact of FHP on balance control and implementing interventions to reduce associated risks.

Key words: centre of pressure, postural stability, forward head posture, balance control, smartphone texting

Introduction

Smartphone usage significantly impacts posture, particularly in the neck and trunk regions, leading to musculoskeletal discomfort and altered spinal kinematics. Studies have shown that smartphone addiction correlates with compromised head posture, decreased muscle endurance, and changes in neck alignment [1]. Additionally, smartphone users exhibit increased neck flexion, head flexion, and forward head posture (FHP), exceeding recommended safe angles [2, 3]. Furthermore, prolonged smartphone use is associated with neck pain, discomfort, and poor sitting neck posture [4]. This altered posture can affect the body's centre of pressure (CoP) distribution and stabilographic variables, potentially impacting gait parameters [5]. Therefore, it is crucial to raise awareness about the ergonomic risks posed by excessive smartphone usage to mitigate musculoskeletal issues and maintain overall postural health.

Advancements in smartphone technology have led to the integration of dual-task demands, such as standing, walking, and texting or talking simultaneously [6]. Research indicates that dual-tasking with smartphones can significantly impact gait parameters, cognitive performance, and stability [7]. Studies have shown that dual-tasking while using smartphones can increase the risk of falls due to reduced situational awareness and cognitive load [8], emphasizing the need for further examination to enhance fall-prevention interventions and community participation. Understanding the effects of smartphone-related dual-tasking on gait and cognitive performance is crucial for improving safety measures and designing interventions to mitigate potential risks associated with these everyday activities.

Maintaining a straight posture while using smartphones affects postural control and motor performance [5, 9, 10]. Research indicates that older individuals experience a more significant decline in postural sta-

Correspondence address: Weerasak Tapanya, Department of Physical Therapy, School of Allied Health Sciences, University of Phayao, 19/2 Maeka Subdistrict, Muang District, Phayao 56000, Thailand, e-mail: weerasak.ta@up.ac.th; <https://orcid.org/0000-0003-2107-0326>

Received: May 13, 2024

Accepted for publication: August 11, 2024

Citation: Tapanya W, Sangkarit N. Effects of forward head posture on the centre of pressure sway during dual-task balancing with smartphone engagement. *Hum Mov.* 2024;25(3):31–42; <https://doi.org/10.5114/hm/192231>.

bility from smartphone use compared to younger individuals, with cognitive function deficits contributing to decreased performance during dual-tasking [11]. The complexity of postural control is further illustrated by analyzing the CoP during cognitive dual-tasks, demonstrating reduced efficacy in postural control [10]. These findings underscore the importance of considering how smartphone use impacts postural control and motor performance in everyday tasks.

Texting while walking has a more significant detrimental effect on balance, gait, and walking behaviour. Studies indicate that texting while walking leads to a reduction in gait speed, cadence, stride length, and step length, along with increased hip and ankle stiffness, which can increase the risk of fatigue [12]. Additionally, texting has been associated with a higher dual-tasking cost in terms of walking time outdoors compared to indoors, emphasizing the impact of texting on dynamic tasks in different environments [13]. Previous research has shown that texting while walking has a notable impact on balance. This is due to the dynamic nature of walking, which naturally presents challenges. However, there is no existing evidence to demonstrate that texting on a smartphone affects static balance, such as when standing on one leg. Exploring the repercussions of multitasking could greatly enrich the evidence base, aiding in understanding the mechanisms and effects of smartphone use on health, such as texting on a smartphone while balancing.

FHP significantly impacts postural stability and balance. Individuals with severe FHP exhibit impaired dynamic postural stability in the sagittal plane, which emphasizes the importance of assessing dynamic balance in FHP participants [14]. Therefore, based on the hypothesis, individuals with FHP during smartphone texting are likely to experience greater alterations in CoP during standing, particularly due to the dual-task demands, increasing their risk compared to those with normal posture. However, there has been no study comparing the effects of smartphone texting while balancing on one leg between these two groups. Previous research has shown that using a smartphone with one hand vs. two hands affects various variables. Hence, comparing the effects of using a smartphone with one hand vs. two hands on CoP during standing is of interest. The objective of this study is to compare the effects of using a smartphone with one hand vs. two hands on CoP during single-leg standing between individuals with FHP and those with normal posture. This study will offer valuable insights into the health implications of FHP during smartphone use by examining its effects on postural stability and balance, shedding light on

how FHP influences musculoskeletal health and functional abilities. Investigating differences in postural control between one-handed (1H) and two-handed smartphone use (2H) will also help understand how smartphone interactions impact individuals with FHP, potentially guiding ergonomic interventions and rehabilitation programs to mitigate the negative health effects of prolonged smartphone use.

Material and methods

Study design

The present cross-sectional study employed convenience sampling methods to enrol participants, utilizing printed ads and social media platforms for recruitment. A randomized repeated design was used to investigate single-leg standing balance in three conditions: non-smartphone usage (NU; single task), 1H (dual tasks), and 2H (dual tasks), on soft and hard surfaces. The study was conducted at the Physical Therapy Laboratory, Faculty of Allied Health Sciences, University of Phayao, Thailand. CoP sway was assessed with the Nintendo Wii Balance Board (NWBB) for each condition while participants engaged in text typing on a smartphone and performed a single-leg standing test on both surfaces.

Participants

In the research, 57 young and healthy adults were initially part of the study. However, 7 participants were excluded for various reasons: 4 due to excess weight, 2 due to obesity, and 1 for failing the balance test during screening. As a result, the final group was composed of 50 participants aged 18 to 25 years old. All these participants were right-handed and had familiarity with smartphones for at least one year, typically spending about four hours daily on their devices. Participants with a history of trauma or musculoskeletal symptoms in the neck and upper extremities were excluded [15]. None of them had recent discomfort or trauma in the past week. The sample size calculation was conducted using the G*Power software, version 3.1.9.6 for Mac OS, incorporating mean and standard deviation values obtained from a previous investigation on postural sway [16]. An a priori power analysis was conducted to determine the required sample size using an *F*-test for ANOVA with repeated measures and within-between interactions. The partial η^2 was set to 0.05, and the sample size was computed based on the specified alpha level, desired power, and effect size.

A power calculation was carried out with a critical α -value of 0.05, aiming for a strong 90% power level. The results indicated that a minimum of 42 participants was necessary to achieve the predetermined level of statistical significance. Prior to enrollment, all participants underwent validation to confirm the absence of any neurological or musculoskeletal disorders. They also confirmed not engaging in balance-specific training over the last six months. An assessment of limb dominance showed that all participants had dominance in their right leg, matching their dominant leg for kicking the ball [17]. After the first screening process, the participants were divided into two groups based on their craniocervical (CV) angle: a control group ($n = 25$) with an angle $\geq 48^\circ$ and an FHP group ($n = 25$) with an angle $< 48^\circ$ [18]. Demographic characteristics and anthropometric data are presented in Table 1. The demographic and anthropometric variables did not show any statistically significant differences ($p > 0.05$) between the control and FHP groups, except for the CV angle. The normal group presented a significantly higher CV angle (mean \pm SD: 52.98 ± 3.54) than the FHP group (44.66 ± 2.76 ; $p < 0.001$).

CV angle

CV angle is measured using a method that involves photometry [19]. The CV angle is defined as the angle formed between a horizontal line through the seventh cervical vertebra (C7) and a line drawn from the tragus of the ear to the C7 vertebra. This angle is used to assess the alignment of the head and neck, particularly in relation to FHP. A smaller CV angle ($< 48^\circ$) indicates a more pronounced FHP, while a larger angle ($\geq 48^\circ$) indicates a more neutral head position [18]. A camera is placed 1.5 m away and adjusted to the participant's shoulder level [19]. To ensure accuracy, markers are placed at C7 and the ear tragus. Participants are asked

to stand straight, facing forward, for a lateral view photo [19]. The digital camera used was a Fujifilm (X-T100 model; 24.2 million pixels; Fujifilm Corp., Japan). The images were then analyzed using Kinovea software, drawing lines between the tragus midpoint and C7 spinous process, as well as a horizontal line through C7. A CV angle below 48° indicates FHP. CV angle measurements demonstrate a high level of reliability, with a consistent test-retest reliability of 0.85–0.86 and an inter-examiner intraclass correlation coefficient of 0.88–0.89 [20].

Experimental protocol

Written consent was obtained from all participants before the study began. Participants were randomly allocated the testing sequence of six different conditions: NU, 1H (dual tasks), and 2H (dual tasks), each on soft and hard surfaces (as illustrated in Figure 1). This allocation process involved basic randomization to avoid any potential biases. During evaluations for NU, participants were asked to assess their balance on one leg while folding both hands across their chest. For smartphone usage positions, participants were instructed to maintain a neck angle of 0 – 15° flexion, in line with ergonomic recommendations. Shoulder angles, smartphone height, and distance were customized according to each participant's usual smartphone habits to ensure a comfortable experience. Participants were allowed 3 min to acclimate to the measuring procedure [15]. In each smartphone usage scenario, participants were required to perform a single-leg standing test on their dominant leg while typing the English alphabet (A–Z) with their thumb, securely holding the device in their hand. Each session lasted 1 min, with a 2-min break between trials.

The single-leg standing evaluation began with participants standing with both feet on the surface. They

Table 1. Demographic and anthropometric characteristics of the participants ($n = 50$)

Variables	Normal group ($n = 25$) mean \pm SD	FHP group ($n = 25$) mean \pm SD	p -value
Gender (males / females)	6(24%) / 19(76%)	9(36%) / 19(76%)	0.365
Age	20.32 ± 1.07	20.48 ± 0.96	0.581
Weight	53.66 ± 6.71	56.60 ± 11.12	0.264
Height	162.48 ± 7.26	163.88 ± 9.07	0.550
BMI	20.24 ± 1.61	20.87 ± 2.09	0.239
CV angle (degree)	52.98 ± 3.54	44.66 ± 2.76	$< 0.001^{**}$
Total experience time of smartphone use (years)	8.19 ± 2.05	8.34 ± 2.09	0.796
Daily duration of smartphone use (hours)	6.66 ± 1.39	6.44 ± 1.55	0.612

BMI – body mass index, CV – craniocervical, ** indicates a significant difference between conditions at p -value < 0.01



Figure 1. The smartphone texting task performed during single-leg standing under various conditions

then transferred their weight to their dominant leg as instructed, maintaining control of the smartphone and ensuring that the non-dominant knee was flexed at a 90° angle [17].

NWBB for assessment of postural sway during quiet standing

CoP was assessed using the NWBB (Nintendo, Kyoto, Japan). The NWBB is made from a plastic composite material and has four load sensors at each corner to measure vertical forces. It uses a Bluetooth connection to link with a laptop computer, which is controlled by customized LabVIEW software (National Instruments, Austin, TX, USA). Data are transferred via Windows-based software designed for Bluetooth. The data are sampled at 30 Hz, consistent with previous studies (10–50 Hz), and a 10 Hz cut-off filter is applied post-sampling [21, 22]. Before each test, the NWBB was calibrated with a 20.4 kg weight [23]. CoP variables such as path length sway, anteroposterior (AP) sway amplitude, and mediolateral (ML) sway amplitude were measured in centimetres.

Statistical analysis

Participant demographic data were initially summarized using descriptive statistics such as means and standard deviations. A Shapiro–Wilk test was used to assess the normal distribution of variables. The Chi-square test was used to examine the differences in gender between groups. A two-way mixed ANOVA was

performed to evaluate the interaction effects between conditions (NU, 1H, and 2H) and groups (FHP and normal) on CoP sway variables in all directions. Independent *t*-tests were conducted to compare CoP sway variables between groups under different test conditions. Within-group comparisons across conditions were performed using one-way repeated measures ANOVA, followed by Bonferroni post hoc analysis to identify pairwise differences. Data analysis was conducted using SPSS software version 26.0 (IBM SPSS Statistics, SPSS Inc., Chicago, IL, USA), with a predetermined significance level (α) set at 0.05.

Results

The interaction effects (group \times conditions) on CoP sway were significant for all variables, including path length sway, AP sway amplitude, and ML sway amplitude, on both soft and hard surfaces (partial $\eta^2 = 0.966$ to 0.988 , $p < 0.001$). The between-group (normal vs. FHP) interaction effects were significant for all variables on both soft and hard surfaces (partial $\eta^2 = 0.187$ to 0.373 , $p < 0.01$). Additionally, the within-group (NU vs. 1H vs. 2H) interaction effects were significant for all variables on both soft and hard surfaces (partial $\eta^2 = 0.379$ to 0.752 , $p < 0.001$). Post hoc tests revealed significant differences between NU vs. 1H and NU vs. 2H, but no significant difference was found between 1H vs. 2H, as shown in Table 2.

When comparing the normal and FHP groups, it was observed that the FHP group exhibited greater CoP sway length in all directions, including path length

Table 2. Two-way, mixed-effects analysis of variance of covariance with repeated measures (2 groups × 3 conditions), between-group interaction effect (normal vs. FHP), and within-group interaction effect (NU vs. 1H vs. 2H).

Variable	Testing condition	Mean ± standard error			Mean differences (between groups)	Between-group interaction effect <i>p</i> -value (partial η^2)	Group × condition interaction effect <i>p</i> -value (partial η^2)
		all groups	normal group	FHP group			
Path length sway (cm)	NU/HS	91.96 ± 2.80			-23.30 ± 5.54	< 0.001** (0.269)	< 0.001** (0.970)
	1H/HS	119.63 ± 3.96 ^a					
	2H/HS	117.98 ± 3.10 ^b					
	within group interaction effect <i>p</i> -value (partial η^2)	< 0.001** (0.606)					
Path length sway (cm)	NU/SS	108.32 ± 3.21			-35.94 ± 7.76	< 0.001** (0.309)	< 0.001** (0.966)
	1H/SS	157.39 ± 4.95 ^a					
	2H/SS	162.04 ± 5.17 ^b					
	within group interaction effect <i>p</i> -value (partial η^2)	< 0.001** (0.752)					
AP sway amplitude (cm)	NU/HS	3.43 ± 0.12			-0.75 ± 0.14	< 0.001** (0.373)	< 0.001** (0.984)
	1H/HS	3.86 ± 0.10 ^a					
	2H/HS	4.11 ± 0.08 ^b					
	within-group interaction effect <i>p</i> -value (partial η^2)	< 0.001** (0.379)					
AP sway amplitude (cm)	NU/SS	4.02 ± 0.10			-0.68 ± 0.21	0.002** (0.187)	< 0.001** (0.978)
	1H/SS	4.88 ± 0.12 ^a					
	2H/SS	5.23 ± 0.21 ^b					
	within-group interaction effect <i>p</i> -value (partial η^2)	< 0.001** (0.485)					
ML sway amplitude (cm)	NU/HS	2.54 ± 0.07			-0.40 ± 0.11	< 0.001** (0.205)	< 0.001** (0.984)
	1H/HS	3.32 ± 0.08 ^a					
	2H/HS	3.40 ± 0.08 ^b					
	within-group interaction effect <i>p</i> -value (partial η^2)	< 0.001** (0.667)					
ML sway amplitude (cm)	NU/SS	2.90 ± 0.07			-0.46 ± 0.11	< 0.001** (0.263)	< 0.001** (0.988)
	1H/SS	3.79 ± 0.07 ^a					
	2H/SS	3.91 ± 0.09 ^b					
	within-group interaction effect <i>p</i> -value (partial η^2)	< 0.001** (0.706)					

NU/HS – non-usage on hard surface, 1H/HS – 1 hand on hard surface, 2H/HS – 2 hands on hard surface

NU/SS – non-usage on soft surface, 1H/SS – 1 hand on soft surface, 2H/SS – 2 hands on soft surface

* significant difference at *p* < 0.05, ** significant difference at *p* < 0.001

^a significant difference between NU and 1H, ^b significant difference between NU and 2H

Table 3. Comparison of CoP sway variables in each direction between FHP and normal groups, on both soft and hard surfaces ($n = 50$)

Variable	Testing condition	Mean \pm SD		Mean differences (between groups)	Between-group p -value (95% CI)
		normal group	FHP group		
Path length sway (cm)	NU/HS	83.16 \pm 20.06	100.76 \pm 19.49	-17.60 \pm 5.59	0.003** (-28.85 to -6.35)
	1H/HS	107.12 \pm 26.82 ^a	132.14 \pm 29.14 ^a	-25.02 \pm 7.92	0.003** (-40.94 to -9.10)
	2H/HS	104.35 \pm 20.10 ^b	131.62 \pm 23.64 ^b	-27.27 \pm 6.21	< 0.001** (-39.75 to -14.79)
	NU/SS	98.28 \pm 22.15	118.35 \pm 23.18	-20.07 \pm 6.41	0.003** (-32.97 to -7.18)
	1H/SS	139.12 \pm 26.63 ^a	175.66 \pm 41.74 ^a	-36.54 \pm 9.90	< 0.001** (-56.45 to -16.63)
	2H/SS	136.42 \pm 27.72 ^b	187.65 \pm 43.65 ^{b,c}	-51.22 \pm 10.34	< 0.001** (-72.02 to -30.43)
AP sway amplitude (cm)	NU/HS	3.10 \pm 0.66	3.76 \pm 0.97	-0.65 \pm 0.23	0.008** (-1.12 to -0.18)
	1H/HS	3.47 \pm 0.66 ^a	4.25 \pm 0.70 ^a	-0.78 \pm 0.19	< 0.001** (-1.16 to -0.39)
	2H/HS	3.70 \pm 0.55 ^b	4.52 \pm 0.61 ^b	-0.82 \pm 0.16	< 0.001** (-1.15 to -0.49)
	NU/SS	3.82 \pm 0.65	4.23 \pm 0.75	-0.42 \pm 0.20	0.041* (-0.82 to -0.02)
	1H/SS	4.60 \pm 0.85 ^a	5.17 \pm 0.86 ^a	-0.58 \pm 0.24	0.021* (-1.06 to -0.09)
	2H/SS	4.70 \pm 1.13 ^b	5.75 \pm 1.74 ^b	-1.05 \pm 0.42	0.014* (-1.89 to -0.22)
ML sway amplitude (cm)	NU/HS	2.35 \pm 0.47	2.72 \pm 0.50	-0.36 \pm 0.14	0.010* (-0.64 to -0.09)
	1H/HS	3.10 \pm 0.48 ^a	3.53 \pm 0.66 ^a	-0.43 \pm 0.16	0.011* (-0.76 to -0.10)
	2H/HS	3.21 \pm 0.61 ^b	3.60 \pm 0.54 ^b	-0.39 \pm 0.16	0.020* (-0.72 to -0.06)
	NU/SS	2.73 \pm 0.44	3.08 \pm 0.58	-0.35 \pm 0.15	0.020* (-0.64 to -0.06)
	1H/SS	3.55 \pm 0.43 ^a	4.03 \pm 0.60 ^a	-0.48 \pm 0.15	0.002** (-0.77 to -0.18)
	2H/SS	3.63 \pm 0.47 ^b	4.19 \pm 0.76 ^b	-0.56 \pm 0.18	0.003** (-0.92 to -0.20)

NU/HS – non-usage on hard surface, 1H/HS – 1 hand on hard surface, 2H/HS – 2 hands on hard surface


NU/SS – non-usage on soft surface, 1H/SS – 1 hand on soft surface, 2H/SS – 2 hands on soft surface

FHP – forward head posture

For between groups, * significant difference at $p < 0.05$, ** significant difference at $p < 0.001$

For within groups, ^a significant difference between NU and 1H, ^b significant difference between NU and 2H

^c significant difference between 1H and 2H



For between groups, * significant difference at $p < 0.05$, ** significant difference at $p < 0.001$. For within groups, "a" significant difference between NU and 1H, "b" significant difference between NU and 2H, "c" significant difference between 1H and 2H.

Figure 2. Comparative analysis of CoP sway between groups; the FHP group and the normal group, when utilizing a smartphone and maintaining a single-leg stance in different conditions, and within-group between three conditions; NU, 1H, and 2H on hard and soft surfaces.

sway, AP sway amplitude, and ML sway amplitude, across all conditions: NU (single task), 1H (dual tasks), and 2H (dual tasks), on both soft and hard surfaces. These differences were statistically significant ($p < 0.05$), as shown in Table 3 and Figure 2.

Within-group comparisons of smartphone usage revealed that both 1H and 2H smartphone usage led to a significant increase in CoP sway length compared to NU in all directions, on both soft and hard surfaces ($p < 0.05$), in both groups. However, no significant differences were observed between 1H and 2H, except in the FHP group, where 2H resulted in a significantly greater increase in path length sway compared to 1H on soft surfaces ($p < 0.05$). No such difference was observed in the normal group, as shown in Table 3 and Figure 2.

Discussion

This study is dedicated to investigating the complex interplay between smartphone use and postural stability control, with a specific focus on CoP when balancing on a single leg. The primary objective is to explore how this relationship varies between individuals with FHP and those with normal posture. It also aims to investigate the differences within each group based on diverse patterns of smartphone use, including non-usage, one-hand, and two-hand use, on various surfaces. The study aims to uncover the impact of smartphone dependency on the musculoskeletal health of individuals with FHP by analyzing the complex interactions and emphasizing the importance of understanding how smartphones affect posture control in vulnerable groups such as those with FHP due to their widespread use in society. Additionally, the study aims to investigate how specific smartphone usage patterns affect postural stability in individuals with FHP compared to those with normal posture, potentially leading to tailored strategies to mitigate the negative musculoskeletal effects of smartphone use, especially for high-risk groups, and bridging the gap between technology use and musculoskeletal health outcomes to inform clinical practices and public health initiatives promoting healthier smartphone behaviours and reducing hazards of prolonged use, particularly for individuals with FHP.

The findings of this study unequivocally demonstrate that when comparing individuals' attempts to maintain balance on one leg on both hard and soft surfaces, significant differences arise between those using smartphones and those not using smartphones. Specifically, it was observed that smartphone usage

led to increased CoP sway across all directions, including path length, AP sway, and ML sway. Engaging in a dual task while using smartphones, such as walking or maintaining balance, can significantly impact one's stability and increase the risk of falls [8, 9]. When individuals divide their attention between using a smartphone and performing a physical task simultaneously, it can lead to decreased awareness of their surroundings and compromised balance control [9, 24]. Research suggests that dual-task activities involving smartphones can interfere with postural stability by diverting attention away from maintaining proper balance and body alignment [9]. This distraction can disrupt the coordination between sensory inputs, such as visual and proprioceptive cues, which are essential for balance control [25]. As a result, individuals may exhibit altered gait patterns, reduced stability, and an increased likelihood of losing balance or falling. Moreover, prolonged use of smartphones often involves prolonged periods of static posture, such as neck flexion and decreased head stability, which can further exacerbate the risk of falls [26, 27]. Additionally, the repetitive nature of smartphone use may contribute to muscle fatigue and decreased muscle activation, further compromising balance and stability.

The findings of this study indicate that when comparing CoP sway between the FHP and normal posture groups, the FHP group exhibited a more pronounced increase in CoP sway across all conditions. This difference was evident in every direction and on both hard and soft surfaces, regardless of smartphone usage (non-usage, one-hand use, and two-hand use). The increased CoP sway observed in the FHP group, compared to the normal posture group, during smartphone use and single-leg balancing likely results from several interconnected mechanisms. FHP is characterized by a forward position of the head relative to the shoulders, which can lead to altered alignment of the cervical spine and increased muscle tension in the neck and upper back [28, 29]. This musculoskeletal imbalance can compromise the FHP group's ability to maintain proper postural control during challenging tasks such as single-leg balancing. Proper postural control relies on the integration of sensory inputs from the visual, vestibular, and proprioceptive systems [30, 31]. Individuals with FHP may experience disruptions in sensory integration due to changes in head and neck alignment, leading to decreased proprioceptive feedback and impaired balance coordination [32–34]. The altered alignment associated with FHP can affect neuromuscular coordination patterns, particularly in the muscles of the neck, shoulders, and upper back [35]. Dysfunctional

muscle recruitment and coordination may result in inefficient postural adjustments during dynamic tasks such as single-leg balancing, contributing to increased CoP sway [36].

Engaging in dual-task activities, such as using a smartphone while balancing on one leg, imposes additional cognitive demands on individuals [8]. The FHP group may have more difficulty allocating cognitive resources effectively due to their compromised musculoskeletal and sensory systems associated with their posture [37]. This increased cognitive load can further disrupt postural control mechanisms and worsen CoP sway [38]. Individuals with FHP may adopt compensatory strategies to maintain balance, such as relying more on visual feedback or muscle co-contraction. However, these strategies may be less efficient or effective compared to those employed by individuals with normal posture, leading to greater variability in CoP sway.

Furthermore, when comparing within each group across conditions of one-hand and two-hand smartphone use, no significant differences in CoP sway were observed in any direction, except within the FHP group. Specifically, individuals with FHP exhibited an increase in CoP path length sway when using two hands compared to one hand. This discrepancy in CoP sway based on smartphone handling was not evident in the normal posture group, aligning with earlier findings of CoP sway differences between the two groups. Thus, it is emphasized that the FHP group is at higher risk of losing balance while using a smartphone than the normal posture group. The increased CoP sway seen during two-hand smartphone use compared to one-hand use in single-leg balancing among individuals with FHP involves various factors. Using a smartphone with two hands can exacerbate musculoskeletal strain in individuals with FHP, potentially causing increased muscle tension and altered posture that affect balance [39]. Two-hand smartphone interaction may require more cognitive and visual attention than one-hand use, which can detract from balance maintenance [9], especially in those with FHP who may have limited attentional resources for postural control. Individuals with FHP, who may have reduced neck mobility [40], might struggle to visually scan their surroundings or adjust their posture effectively while using a smartphone with two hands, possibly leading to compensatory movements that increase CoP sway during single-leg balancing. These results highlight the need to examine the complex relationship between smartphone use, postural control, and musculoskeletal factors, particularly in vulnerable populations such as those with FHP.

The limitations of this study include the lack of exploration into the mechanisms underlying the effects of smartphone usage on postural stability. Incorporating measures of muscle activity and sensory feedback could provide valuable insights into these mechanisms. Additionally, the absence of longitudinal follow-up hinders understanding of the long-term impacts of smartphone usage on musculoskeletal health. Future studies should consider longitudinal assessments to provide further insight into the effects of smartphone usage over time, thereby informing preventive strategies and interventions.

Conclusions

In conclusion, our study highlights that individuals with FHP face an elevated risk of losing balance during non-smartphone activities, such as single task, compared to those with normal posture. Furthermore, this risk of balance loss becomes more pronounced when engaging in smartphone use while maintaining single-leg standing balance (a dual task), as evidenced by increased CoP sway compared to the normal posture group. Notably, the risk is particularly heightened during two-hand smartphone use. These findings underscore the importance of considering the impact of FHP on balance control, especially in the context of smartphone usage, and emphasize the need for targeted interventions to mitigate these associated risks.

Data availability statement

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

Acknowledgements

We would like to thank all participants who volunteered to participate in this study.

Ethical approval

The research related to human use has complied with all the relevant national regulations and institutional policies, has followed the tenets of the Declaration of Helsinki, and has been approved by the Institutional Review Board at the University of Phayao, Thailand, approved all experimental protocols (approval No.: UP-HEC 1.2/079/66).

Informed consent

Informed consent has been obtained from all individuals included in this study.

Conflict of interest

The authors state no conflict of interest.

Disclosure statement

No author has any financial interest or received any financial benefit from this research.

Funding

This research project received no external funding.

References

- [1] Torkamani MH, Mokhtarinia HR, Vahedi M, Gabel CP. Relationships between cervical sagittal posture, muscle endurance, joint position sense, range of motion and level of smartphone addiction. *BMC Musculoskelet Disord.* 2023;24(1):61; doi: 10.1186/s12891-023-06168-5.
- [2] Chen Y-L, Chen K-H, Cheng Y-C, Chang C-C. Field study of postural characteristics of standing and seated smartphone use. *Int J Environ Res Public Health.* 2022;19(8):4583; doi: 10.3390/ijerph19084583.
- [3] Song D, Kim E, Kwon Y, Kim H, Shin G. Changes in the range of head flexion rotation and neck muscle activity after prolonged use of a smartphone. *Proc Hum Factors Ergon Soc Ann Meet.* 2020;64(1):930–3; doi: 10.1177/1071181320641222.
- [4] Ojha R, Sindhu B, Sen S. Effects of smartphone addiction on sitting neck posture & hand discomfort: a cross sectional study. *Int J Health Sci.* 2022; 6(2):13642–50; doi: 10.53730/ijhs.v6nS2.8595.
- [5] Onofrei RR, Amaricai E, Suciú O, David VL, Rata AL, Hogeá E. Smartphone use and postural balance in healthy young adults. *Int J Environ Res Public Health.* 2020;17(9):3307; doi: 10.3390/ijerph17093307.
- [6] Almajid R, Appiah-Kubi KO, Cipriani D, Goel R. Dual-tasking interference is exacerbated outdoors: a pilot study. *Front Sports Active Living.* 2023;5; doi: 10.3389/fspor.2023.1077362.
- [7] Shahidian H, Begg R, Ackland DC. The influence of cell phone usage on dynamic stability of the body during walking. *J Appl Biomech.* 2022;38(6): 365–72; doi: 10.1123/jab.2021-0374.
- [8] Carneros-Prado D, Dobrescu CC, González I, Fontecha J, Johnson E, Hervás R. Analysis of dual-tasking effect on gait variability while interacting with mobile devices. *Mathematics.* 2023;11(1): 202; doi: 10.3390/math11010202.
- [9] Lino TB, Scarmagnan GS, Sobrinho-Junior SA, Tessari GMF, Gonçalves GH, Pereira HM, Christofolletti G. Impact of using smartphone while walking or standing: a study focused on age and cognition. *Brain Sci.* 2023;13(7):987; doi: 10.3390/brainsci13070987.
- [10] Saraiva M, Fernandes OJ, Vilas-Boas JP, Castro MA. Standing posture in motor and cognitive dual-tasks during smartphone use: linear and non-linear analysis of postural control. *Eur J Investig Health Psychol Educ.* 2022;12(8):1021–33; doi: 10.3390/ejihpe12080073.
- [11] Ozdemir RA, Contreras-Vidal JL, Lee B-C, Paloski WH. Cortical activity modulations underlying age-related performance differences during posture-cognition dual tasking. *Exp Brain Res.* 2016; 234(11):3321–34; doi: 10.1007/s00221-016-4730-5.
- [12] Lu Z, Zhang X, Mao C, Liu T, Li X, Zhu W, Wang C, Sun Y. Effects of mobile phone use on gait and balance control in young adults: a hip-ankle strategy. *Bioengineering.* 2023;10(6):665; doi: 10.3390/bioengineering10060665.
- [13] Sajewicz J, Dziuba-Słonina A. Texting on a smartphone while walking affects gait parameters. *Int J Environ Res Public Health.* 2023;20(5):4590; doi: 10.3390/ijerph20054590.
- [14] Ahmadipoor A, Khademi-Kalantari K, Rezasoltani A, Naimi S-S, Akbarzadeh-Baghbani A. Effect of Forward Head Posture on dynamic balance based on the biodex balance system. *J Biom Phys Eng.* 2022;12(5):543–8; doi: 10.31661/jbpe.v0i0.1912-1036.
- [15] Xie Y, Szeto GP, Dai J, Madeleine P. A comparison of muscle activity in using touchscreen smartphone among young people with and without chronic neck-shoulder pain. *Ergonomics.* 2016; 59(1):61–72; doi: 10.1080/00140139.2015.1056237.
- [16] Abbasi H, Alizadeh MH, Rajabi R, Mohammadi F. Comparison of static and dynamic postural stability between individuals with and without forward head posture. *Phys Treatments.* 2020;10(3): 127–34.
- [17] Tapanya W, Maharan S, Amput P, Sangkarit N, Suwannakul B. The influence of knee extensor and ankle plantar flexor strength on single-leg standing balance in older women. *J Funct Morphol Kinesiol.* 2023;8(2):67; doi: 10.3390/jfmk8020067.
- [18] Salahzadeh Z, Maroufi N, Ahmadi A, Behtash H, Razmjoo A, Gohari M, Parnianpour M. Assessment of forward head posture in females: observational and photogrammetry methods. *J Back Musculoskelet Rehabil.* 2014;27(2):131–9; doi: 10.3233/BMR-130426.

- [19] Fernández-de-las-Peñas C, Alonso-Blanco C, Cuadrado ML, Pareja JA. Forward head posture and neck mobility in chronic tension-type headache: a blinded, controlled study. *Cephalalgia*. 2006;26(3):314–9; doi: 10.1111/j.1468-2982.2005.01042.x.
- [20] Gallego-Izquierdo T, Arroba-Díaz E, García-Ascoz G, Val-Cano MDA, Pecos-Martin D, Cano-de-la-Cuerda R. Psychometric proprieties of a mobile application to measure the craniovertebral angle a validation and reliability study. *Int J Environ Res Public Health*. 2020;17(18):6521; doi: 10.3390/ijerph17186521.
- [21] Haworth J, Stergiou N. Orderliness of visual stimulus motion mediates sensorimotor coordination. *Fron Physiol*. 2018;9:1441; doi: 10.3389/fphys.2018.01441.
- [22] Rand TJ, Mukherjee M. Transitions in persistence of postural dynamics depend on the velocity and structure of postural perturbations. *Exp Brain Res*. 2018;236(5):1491–500; doi: 10.1007/s00221-018-5235-1.
- [23] Meade ZS, Marmelat V, Mukherjee M, Sado T, Takahashi KZ. Comparison of a portable balance board for measures of persistence in postural sway. *J Biomech*. 2020;100:109600; doi: 10.1016/j.jbiomech.2020.109600.
- [24] Luo Y, Grimaldi N, Zheng H, Giang WCW, Hu B. Distraction from smartphones changed pedestrians' walking behaviors in open areas. *Motor Control*. 2023;27(2):275–92; doi: 10.1123/mc.2022-0023.
- [25] Riem LI, Schmit BD, Beardsley SA (eds.). The Effect of Discrete Visual Perturbations on Balance Control during Gait. In: 2020 42nd Annual International Conference of the IEEE Engineering in Medicine & Biology Society (EMBC); 2020 20–24 July 2020; doi: 10.1109/EMBC44109.2020.9176303.
- [26] Dandumahanti BP, Subramaniyam M. Influence of smartphone game play on head flexion angle, muscle activity, and load at C7 among adolescents. *Work*. 2023;75:1413–25; doi: 10.3233/WOR-220422.
- [27] Wah SW, Chatchawan U, Chatprem T, Puntumetakul R. Prevalence of static balance impairment and associated factors of university student smartphone users with subclinical neck pain: cross-sectional study. *Int J Environ Res Public Health*. 2022;19(17):10723; doi: 10.3390/ijerph191710723.
- [28] Poursadegh M, Azghani MR, Chakeri Z, Okhravi SM, Salahzadeh Z. Postures of the head, upper, and lower neck in forward head posture: static and quasi-static analyses. *Middle East J Rehabil Health Stud*. 2023;10(4):e136377; doi: 10.5812/mejrh-136377.
- [29] Suwaidi ASA, Moustafa IM, Kim M, Oakley PA, Harrison DE. A comparison of two forward head posture corrective approaches in elderly with chronic non-specific neck pain: a randomized controlled study. *J Clin Med*. 2023;12(2):542; doi: 10.3390/jcm12020542.
- [30] Shishkin N, Kitov V, Sayenko D, Tomilovskaya E. Sensory organization of postural control after long term space flight. *Front Neural Circuits*. 2023;17:1135434; doi: 10.3389/fncir.2023.1135434.
- [31] Chaudhary S, Saywell N, Taylor D. The differentiation of self-motion from external motion is a prerequisite for postural control: a narrative review of visual-vestibular interaction. *Front Hum Neurosci*. 2022;16:697739; doi: 10.3389/fnhum.2022.697739.
- [32] Ha S-Y, Sung Y-H. A temporary forward head posture decreases function of cervical proprioception. *J Exerc Rehabil*. 2020;16(2):168–74; doi: 10.12965/jer.2040106.053.
- [33] Migliarese S, White E. Review of forward-head posture and vestibular deficits in older adults. *Curr Geriatr Rep*. 2019;8(3):194–201; doi: 10.1007/s13670-019-00292-8.
- [34] Raoufi Z, Sarrafzadeh J, Emrani A, Ghorbanpour A. Interaction between proprioception, forward head posture and neck pain in adult women. *Func Disabil J*. 2020;2:12.
- [35] Kawasaki T, Ohji S, Aizawa J, Sakai T, Hirohata K, Koseki T, Kuruma H, Okawa A, Jinno T. Characteristics of head and neck alignment and function of deep cervical flexor muscles in patients with nonspecific neck pain. *J Bodyw Mov Ther*. 2024;39:565–71; doi: 10.1016/j.jbmt.2024.03.063.
- [36] Kim E-K, Kim S-G. Forward head posture (FHP) angle and plantar pressure resulting from oscillatory stimulation training of the shoulder joint: a randomized controlled trial. *J Back Musculoskelet Rehabil*. 2019;32:37–42; doi: 10.3233/BMR-160748.
- [37] Oh H-J, Song G-B. Effects of neurofeedback training on the brain wave of adults with forward head posture. *J Phys Ther Sci*. 2016;28(10):2938–41; doi: 10.1589/jpts.28.2938.
- [38] Saraiva M, Paszkiel S, Vilas-Boas JP, Castro MA. Influence of cognitive task difficulty in postural control and hemodynamic response in the pre-

- frontal cortex during static postural standing. *Appl Sci.* 2022;12(13):6363; doi: 10.3390/app12136363.
- [39] Tsang SMH, Cheing GLY, Chan JWK. Severity of slouched posture during smartphone use is associated with the musculoskeletal discomfort, daily usage, and school year among adolescents. *Ergonomics.* 2023;66(9):1340–53; doi: 10.1080/00140139.2022.2146208.
- [40] Panihar U, Joshi S. Systematic review and meta-analysis on efficacy of various exercise programs to improve postural parameters, pain and disability in neck. *Comparat Exerc Physiol.* 2023; 19(2):101–10; doi: 10.3920/CEP220074.